

**SUPPLEMENT TO “EXACT CALCULATIONS FOR FALSE
DISCOVERY PROPORTION WITH APPLICATION TO
LEAST FAVORABLE CONFIGURATIONS”**

BY ETIENNE ROQUAIN AND FANNY VILLERS

UPMC University of Paris 6

We provide some proofs for the paper (1). We use the notation
and the equation numbering defined in (1).

1. Proof of (31)-(1). To prove (31)-(1), remark that we may assumed that $G(x) = x$ up to replace $(t_j)_j$ by $(G(t_j))_j$ (because G is continuous increasing). Next, assume $k < k' < m$ and denote $L(r) = \sum_{i=1}^m \mathbf{1}\{p_i \leq t_r\}$. By definition of a step-down procedure,

$$\begin{aligned}
 & \mathbb{P} [|\text{SD}(\mathbf{t})| = k, |\text{SD}(\mathbf{t}')| = k'] \\
 &= \mathbb{P}[\forall j \leq k, L(j) \geq j, L(k+1) = k, \\
 & \quad \forall j, k+1 \leq j \leq k', L(j+1) \geq j, L(k'+2) = k'] \\
 &= \binom{m}{k} \binom{m-k}{k'-k} \mathbb{P}[\forall j \leq k, L(j) \geq j, \forall j, k+1 \leq j \leq k', L(j+1) \geq j, \\
 & \quad p_1, \dots, p_k \leq t_{k+1} < p_{k+1}, \dots, p_{k'} \leq t_{k'+2} < p_{k'+1}, \dots, p_m] \\
 &= \binom{m}{k} \binom{m-k}{k'-k} \mathbb{P} \left[p_1, \dots, p_k \leq t_{k+1} < p_{k+1}, \dots, p_{k'} \leq t_{k'+2} < p_{k'+1}, \dots, p_m, \right. \\
 & \quad \left. \forall j \leq k, \sum_{i=1}^k \mathbf{1}\{p_i \leq t_j\} \geq j, \forall j, k+1 \leq j \leq k', \sum_{i=k+1}^{k'} \mathbf{1}\{p_i \leq t_{j+1}\} \geq j-k \right] \\
 &= \binom{m}{k} \binom{m-k}{k'-k} \mathbb{P} \left[p_{(1)} \leq t_1, \dots, p_{(k)} \leq t_k \right] (1 - t_{k'+2})^{m-k'} \\
 & \quad \times \mathbb{P} \left[t_{k+1} < p_{(1)} \leq t_{k+2}, \dots, p_{(k'-k)} \leq t_{k'+1} \right],
 \end{aligned}$$

AMS 2000 subject classifications: Primary 62J15; secondary 62G10; tertiary 60C05

Keywords and phrases: False discovery rate, false discovery proportion, multiple testing, least favorable configuration, power, equicorrelated multivariate normal distribution, step-up, step-down

where we used that the p -values are i.i.d. Simple computations give that

$$\begin{aligned} & \mathbb{P} \left[t_{k+1} < p_{(1)} \leq t_{k+2}, \dots, p_{(k'-k)} \leq t_{k'+1} \right] \\ &= \mathbb{P} \left[p_{(1)} \leq t_{k+2} - t_{k+1}, \dots, p_{(k'-k)} \leq t_{k'+1} - t_{k+1} \right]. \end{aligned}$$

This leads to (31)-(1). The cases $k < k' = m$ and $k = k'$ are similar.

2. Proof of (16)-(1), (17)-(1) and (18)-(1). Let us state the following general expression: for $1 \leq j \leq k \leq m$,

(1)

$$\mathbb{P}[|\mathcal{H}_0(H) \cap \text{SD}(\mathbf{t})| = j \mid |\text{SD}(\mathbf{t})| = k] = \binom{k}{j} \pi_0^j \pi_1^{k-j} \frac{\Psi_{k,j,F_1}(t_1, \dots, t_k)}{\Psi_k(G(t_1), \dots, G(t_k))},$$

where we let $\Psi_{k,j,F_1}(t_1, \dots, t_k) = \mathbb{P}[U_{(1)} \leq t_1, \dots, U_{(k)} \leq t_k]$, in which the probability is taken over $(U_i)_{1 \leq i \leq k}$ such that $(U_i)_{1 \leq i \leq j}$ are i.i.d. uniform, independently of $(U_i)_{j+1 \leq i \leq k}$ i.i.d. of c.d.f. F_1 .

Applying (1) in the particular cases $j = k$, $j = 0$ and $F_1 = 1$ leads to (16)-(1), (17)-(1) and (18)-(1), respectively. To state (1), we use that $(H_i, p_i)_i$ is a i.i.d. sequence:

$$\begin{aligned} & \mathbb{P}[|\mathcal{H}_0(H) \cap \text{SD}(\mathbf{t})| = j, |\text{SD}(\mathbf{t})| = k] \\ &= \binom{k}{j} \binom{m}{k} \mathbb{P}[\text{SD}(\mathbf{t}) = \{1, \dots, k\}, H_1 = \dots = H_j = 0, H_{j+1} = \dots = H_k = 1] \\ &= \binom{k}{j} \binom{m}{k} \mathbb{P} \left[\forall \ell \leq k, \sum_{i=1}^k \mathbf{1}\{p_i \leq t_\ell\} \geq \ell, \forall i \geq k+1, p_i > t_{k+1}, \right. \\ & \quad \left. H_1 = \dots = H_j = 0, H_{j+1} = \dots = H_k = 1 \right] \\ &= \binom{k}{j} \pi_0^j \pi_1^{k-j} \mathbb{P} \left[\forall \ell \leq k, \sum_{i=1}^k \mathbf{1}\{p_i \leq t_\ell\} \geq \ell \mid H_1 = \dots = H_j = 0, \right. \\ & \quad \left. H_{j+1} = \dots = H_k = 1 \right] \binom{m}{k} (1 - G(t_{k+1}))^{m-k}, \end{aligned}$$

which leads to (1) by applying (19)-(1).

3. Proof of Proposition 3.8 of (1). We focus on the step-up case, when $\rho \in (-1, 1)$ and $m_0 = 1$ (the remaining cases are left to the reader). Without loss of generality, we may assume that the first coordinate corresponds to the true null, that is, $H = (0, 1)$. In this context, the FDP takes

one of the three values: $0, \frac{1}{2}, 1$, according to the location of the test statistics $Y_i = X_i + \mu H_i$ with respect to the critical values z_1 and z_2 . From the definition of a step-up procedure, we may define the region $\mathcal{D}_i = \{(y_1, y_2) \in \mathbb{R}^2 \mid \text{FDP}(y_1, y_2) = i/2\}$, for $i \in \{1, 2\}$, where $\text{FDP}(y_1, y_2)$ denotes the FDP of $\text{SU}(\mathbf{t})$ taken in the p -values $\mathbf{p} = (\bar{\Phi}(y_1), \bar{\Phi}(y_2))$. The regions \mathcal{D}_i are represented on Figure 1(a). Next, since (Y_1, Y_2) follows the EMN model, we may write for $i \in \{1, 2\}$,

$$\begin{aligned} \mathbb{P}(\text{FDP}(Y_1, Y_2) = i/2) &= (2\pi)^{-1} (1 - \rho^2)^{-1/2} \int_{\mathcal{D}_i} \exp \left\{ -\frac{(y_1 - \rho(y_2 - \mu))^2}{2(1 - \rho^2)} - \frac{1}{2}(y_2 - \mu)^2 \right\} dy_1 dy_2 \\ &= (2\pi)^{-1} \int_{\tilde{\mathcal{D}}_i} \exp\{-(u^2 + v^2)/2\} dudv, \end{aligned}$$

by using the substitution $u = (1 - \rho^2)^{-1/2}(y_1 - \rho(y_2 - \mu))$ and $v = y_2 - \mu$, and where the resulting integration domain $\tilde{\mathcal{D}}_i$ is represented on Figure 1(b). Therefore, we obtain

$$\mathbb{P}(\text{FDP} = 1/2) = (2\pi)^{-1/2} \int_{z_1 - \mu}^{\infty} \exp\{-v^2/2\} \bar{\Phi} \left((z_1 - \rho v) / \sqrt{1 - \rho^2} \right) dv,$$

and

$$\mathbb{P}(\text{FDP} = 1) = (2\pi)^{-1/2} \int_{-\infty}^{z_1 - \mu} \exp\{-v^2/2\} \bar{\Phi} \left((z_2 - \rho v) / \sqrt{1 - \rho^2} \right) dv,$$

and the final expression results by using the substitution $w = \bar{\Phi}(v)$.

4. Proof that the linear threshold satisfies condition (A)-(1).

Let us prove that the threshold $t_j = \alpha j/m$, $1 \leq j \leq m$, satisfies (A)-(1). For any $m \geq 2$ and $1 \leq k \leq m$, let us denote

$$S_{m,k} = \sum_{i=0}^{m-k} \frac{k}{k+i} \tilde{\mathcal{D}}_{m-k} \left(\left(\frac{\alpha j/m}{1 - \alpha k/m} \right)_{1 \leq j \leq m-k}, i \right)$$

and prove that $k \in \{1, \dots, m\} \mapsto S_{m,k}$ is nondecreasing. Letting $a_{m,k} = \frac{k}{m} \left(1 - \alpha \frac{m-k}{m-\alpha k} \right)$ (increasing in k) and $b_{m,k} = \frac{m-\alpha}{m} \frac{m-k}{m-\alpha k}$ (decreasing in k), we may prove the following recursion (see the proof below):

$$(2) \quad S_{m,k} = a_{m,k} + b_{m,k} S_{m-1,k}.$$




FIG 1. Top: \mathcal{D}_i . Bottom: $\tilde{\mathcal{D}}_i$. For $i = 1$ (double cross-hatch) and $i = 2$ (simple cross-hatch). Graph for $\rho < 0$.

Using (2), we can state that $k \in \{1, \dots, m\} \mapsto S_{m,k}$ is nondecreasing by a recurrence on $m \geq 2$: the property is obviously true for $m = 2$. Assuming the property true for $m - 1$, we obtain for any $2 \leq k \leq m - 1$,

$$\begin{aligned} S_{m,k} - S_{m,k-1} &= a_{m,k} - a_{m,k-1} + b_{m,k}S_{m-1,k} - b_{m,k-1}S_{m-1,k-1} \\ &= a_{m,k} - a_{m,k-1} + (b_{m,k} - b_{m,k-1})S_{m-1,k} \\ &\quad + b_{m,k-1}(S_{m-1,k} - S_{m-1,k-1}) \\ &\geq (a_{m,k} + b_{m,k}) - (a_{m,k-1} + b_{m,k-1}), \end{aligned}$$

because $S_{m-1,k} \leq 1$. Hence, since $a_{m,k} + b_{m,k} = 1 - \frac{\alpha}{m} \frac{m-k}{m-\alpha k}$, the quantity $a_{m,k} + b_{m,k}$ is increasing in k and $S_{m,k} - S_{m,k-1} \geq 0$. Also, we obviously have $S_{m,m} = 1 \geq S_{m,m-1}$, and the recurrence is completed.

We now finally state (2). Let for $1 \leq j \leq m - k$, $t_j = \frac{\alpha j/m}{1 - \alpha k/m}$ and $t'_j = 1 - t_{m-k-j+1}$, so that

$$S_{m,k} = \mathbb{E}_0 \left[\frac{k}{k + |\text{SD}((t_j)_{1 \leq j \leq m-k})|} \right] = \mathbb{E}_0 \left[\frac{k}{m - |\text{SU}((t'_j)_{1 \leq j \leq m-k})|} \right],$$

where \mathbb{E}_0 denotes the expectation with respect to i.i.d. uniform p -values. Hence, denoting $t'_j = \beta + j\gamma$ with $\beta = \frac{m-m\alpha-\alpha}{m-\alpha k}$ and $\gamma = \frac{\alpha}{m-\alpha k}$, we obtain

$$\begin{aligned} \frac{S_{m,k}}{k} &= \frac{1}{m} + \frac{1}{m} \mathbb{E}_0 \left[\frac{|\text{SU}((t'_j)_{1 \leq j \leq m-k})|}{m - |\text{SU}((t'_j)_{i \leq j \leq m-k})|} \right] \\ &= \frac{1}{m} + \frac{1}{m} \sum_{j=1}^{m-k} \frac{j}{m-j} \binom{m-k}{j} (t'_j)^j \Psi_{m-k-j}(1 - t'_{m-k}, \dots, 1 - t'_{j+1}) \\ &= \frac{1}{m} + \frac{m-k}{m} \sum_{j=1}^{m-k} \frac{t'_j}{m-j} \binom{m-k-1}{j-1} (t'_j)^{j-1} \\ &\quad \times \Psi_{m-k-j}(1 - t'_{m-k}, \dots, 1 - t'_{j+1}). \end{aligned}$$

Therefore,

$$\begin{aligned} \frac{S_{m,k}}{k} &= \frac{1}{m} - \gamma \frac{m-k}{m} + \frac{m-k}{m} (\beta + m\gamma) \\ &\quad \times \sum_{j=0}^{m-1-k} \frac{(t'_{j+1})^j}{m-1-j} \binom{m-1-k}{j} \Psi_{m-1-k-j}(1 - t'_{m-k}, \dots, 1 - t'_{j+2}) \\ &= \frac{1}{m} - \gamma \frac{m-k}{m} + \frac{m-k}{m} (\beta + m\gamma) \mathbb{E}_0 \left[\frac{1}{m-1 - |\text{SU}((t'_{j+1})_{1 \leq j \leq m-1-k})|} \right] \\ &= \frac{1}{m} - \gamma \frac{m-k}{m} + \frac{m-k}{m} (\beta + m\gamma) \frac{S_{m-1,k}}{k}, \end{aligned}$$

and the recursion (2) is proved.

5. Proof of (29)-(1). For any $0 \leq i \leq m-1$, let us denote

$$u_i = \mathbb{E}_0[(|\text{SU}((t_j)_{i+1 \leq j \leq m})| + i + 1)^{-1}],$$

(where \mathbb{E}_0 denotes the expectation with respect to i.i.d. uniform p -values) and $u_m = 1/(m+1)$, $u_{m+1} = 0$, so that u_0 equals the quantity in (29)-(1). We may prove the following recursion relation: for any $0 \leq i \leq m$,

$$(3) \quad (i+1)u_i = (1 - (m-i)\gamma) - (m-i)(\beta - \gamma)u_{i+1}.$$

Expression (3) is proved as follows: for $i < m$,

$$\begin{aligned} (i+1)u_i &= 1 - \mathbb{E}_0 \left[\frac{|\text{SU}((t_j)_{i+1 \leq j \leq m})|}{|\text{SU}((t_j)_{i+1 \leq j \leq m})| + i + 1} \right] \\ &= 1 - \sum_{k=1}^{m-i} \frac{k}{k+i+1} \binom{m-i}{k} (t_{k+i})^k \Psi_{m-i-k}(1-t_m, \dots, 1-t_{k+i+1}) \\ &= 1 - (m-i) \sum_{k=1}^{m-i} \frac{t_{k+i}}{k+i+1} \binom{m-i-1}{k-1} (t_{k+i})^{k-1} \\ &\quad \times \Psi_{m-i-k}(1-t_m, \dots, 1-t_{k+i+1}) \\ &= 1 - (m-i) \sum_{k=0}^{m-(i+1)} \left(\gamma + \frac{\beta - \gamma}{k + (i+1) + 1} \right) \binom{m-(i+1)}{k} (t_{k+(i+1)})^k \\ &\quad \times \Psi_{m-(i+1)-k}(1-t_m, \dots, 1-t_{k+(i+1)+1}). \end{aligned}$$

Next, we obtain that the solution of the recursion (3) is given by

$$u_i = \sum_{j=0}^{m-i} \frac{1 - (m-(i+j))\gamma}{m-(i+j)} \frac{(m-i) \times \dots \times (m-(i+j))}{(i+1) \times \dots \times (i+j+1)} (\gamma - \beta)^j,$$

which leads to $u_0 = \sum_{j=0}^m \frac{1-(m-j)\gamma}{j+1} \binom{m}{j} (\gamma - \beta)^j$ and (29)-(1) results.

6. Proofs of the lemmas of Section 7 in (1).

6.1. *Proof of Lemma 7.1 of (1).* First note that since \hat{k} is nondecreasing in each coordinate of (p_1, \dots, p_m) , we always have $\hat{k} \leq \hat{k}'_{(\ell)} + \ell$. Second, since $p_{(k)} \leq t_k$ is equivalent to $|\{1 \leq j \leq m \mid p_j \leq t_k\}| \geq k$, the rejection number of $\text{SU}(\mathbf{t})$ can be defined as $\hat{k} = \max\{k \in \{0, 1, \dots, m\} \mid |\{1 \leq j \leq m \mid p_j \leq t_k\}| \geq k\}$. Hence, $\forall 1 \leq i \leq \ell$, $p_i \leq t_{\hat{k}'_{(\ell)} + \ell}$ is equivalent to $|\{1 \leq j \leq m \mid p_j \leq t_{\hat{k}'_{(\ell)} + \ell}\}| \geq \ell + |\{\ell + 1 \leq j \leq m \mid p_j \leq t_{\hat{k}'_{(\ell)} + \ell}\}|$ which is

equivalent to $|\{1 \leq j \leq m \mid p_j \leq t_{\hat{k}'_{(\ell)}+\ell}\}| \geq \ell + \hat{k}'_{(\ell)}$, by definition of $\hat{k}'_{(\ell)}$. As a consequence, since \hat{k} is a maximum and since $\hat{k} \leq \hat{k}'_{(\ell)} + \ell$, the latter is equivalent to $\hat{k}'_{(\ell)} + \ell = \hat{k}$. This establishes the second equivalence. The first equivalence easily comes from the second equivalence and using that $t_{\hat{k}} \leq t_{\hat{k}'_{(\ell)}+\ell}$ because $(t_k)_k$ is a nondecreasing sequence.

6.2. *Proof of Lemma 7.2 of (1).* Similarly to the step-up case, the rejection number of $\text{SD}(\mathbf{t})$ can be defined as $\tilde{k} = \max\{k \in \{0, 1, \dots, m\} \mid \forall k' \leq k, |\{1 \leq j \leq m \mid p_j \leq t_{k'}\}| \geq k'\}$. Also remark that we always have $\tilde{k}'_{(1)} + 1 \geq \tilde{k}$ and, by definition of \tilde{k} , for any j we have $p_j \leq t_{\tilde{k}} \Leftrightarrow p_j \leq t_{\tilde{k}+1}$. First prove that $p_1 \leq t_{\tilde{k}} \Leftrightarrow \tilde{k} = \tilde{k}'_{(1)} + 1$: using the definitions of \tilde{k} and $\tilde{k}'_{(1)}$ we obtain $p_1 \leq t_{\tilde{k}} \Leftrightarrow |\{2 \leq j \leq m \mid p_j \leq t_{\tilde{k}+1}\}| < \tilde{k} \Leftrightarrow \tilde{k}'_{(1)} < \tilde{k} \Leftrightarrow \tilde{k}'_{(1)} + 1 = \tilde{k}$. Second, we prove $p_1 > t_{\tilde{k}} \Leftrightarrow p_1 > t_{\tilde{k}+1}$: since we obviously have $\tilde{k} \geq \tilde{k}'_{(1)}$, we get $p_1 > t_{\tilde{k}} \Rightarrow p_1 > t_{\tilde{k}+1} \Rightarrow p_1 > t_{\tilde{k}'_{(1)}+1}$. Conversely, if $p_1 > t_{\tilde{k}'_{(1)}+1}$, we get $|\{1 \leq j \leq m \mid p_j \leq t_{\tilde{k}'_{(1)}+1}\}| = |\{2 \leq j \leq m \mid p_j \leq t_{\tilde{k}'_{(1)}+1}\}| < \tilde{k}'_{(1)} + 1$ (by definition of $\tilde{k}'_{(1)}$), hence $\tilde{k}'_{(1)} + 1 > \tilde{k}$ (by definition of \tilde{k}), which implies $\tilde{k}'_{(1)} = \tilde{k}$, thus $p_1 > t_{\tilde{k}+1}$ and finally $p_1 > t_{\tilde{k}}$.

References.

- [1] ROQUAIN, E. and VILLERS, F. (2011). Exact calculations for false discovery proportion with application to least favorable configurations. *Ann. Statist.* **39** 584–612.

UPMC UNIV PARIS 6, LPMA,
4, PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE
E-MAIL: etienne.roquain@upmc.fr

UPMC UNIV PARIS 6, LPMA,
4, PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE
E-MAIL: fanny.villers@upmc.fr